



HAL
open science

Algebraicity of foliations on complex projective manifolds, applications

Frederic Bruno Campana

► **To cite this version:**

Frederic Bruno Campana. Algebraicity of foliations on complex projective manifolds, applications. 2021. hal-03475769

HAL Id: hal-03475769

<https://univ-poitiers.hal.science/hal-03475769>

Preprint submitted on 11 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ALGEBRAICITY OF FOLIATIONS ON COMPLEX PROJECTIVE MANIFOLDS, APPLICATIONS

FRÉDÉRIC CAMPANA

CONTENTS

1. Algebraicity criterion: statement	2
2. Proof of the algebraicity criterion.	5
3. Pseudoeffectivity and movable classes.	6
4. Harder-Narasimhan filtrations and pseudo-effectivity.	8
5. Pseudo-effectivity of relative canonical bundles.	11
6. Rational curves and non-pseudoeffectivity of the canonical/cotangent bundles.	13
7. Birational stability of the cotangent bundle	15
8. Shafarevich-Viehweg conjecture.	17
9. Numerically trivial foliations.	18
10. The Beauville-Bogomolov-Yau decomposition	19
11. Some questions	21
11.1. Pseudoeffectivity of determinants.	21
11.2. Variations on Abundance.	22
11.3. Special manifolds.	23
References	24

ABSTRACT. This is an expository text, originally intended for the ANR ‘Hodgefun’ workshop, twice reported, organised at Florence, villa Finaly, by B. Klingler.

We show that holomorphic foliations on complex projective manifolds have algebraic leaves under a certain positivity property: the ‘non pseudoeffectivity’ of their duals. This permits to construct certain rational fibrations with fibres either rationally connected, or with trivial canonical bundle, of central importance in birational geometry. A considerable extension of the range of applicability is due to the fact that this positivity is preserved by the tensor powers of the tangent bundle. The results presented here are extracted from [10], which is inspired by the former results ([28], [6], [11]). In order to make things as simple as possible, we present here only the projective versions of these results, although most of them can be easily extended to the logarithmic or ‘orbifold’ context.

1. ALGEBRAICITY CRITERION: STATEMENT

In all the text, X will denote an n -dimensional connected complex projective manifold, A will be an ample line bundle on X . A Zariski open subset of X is said to be ‘big’ if its complement has codimension at least 2 in X .

Let $\mathcal{F} \subset TX$, the holomorphic tangent bundle of X , be a coherent saturated subsheaf of rank $r \leq n$.

Recall that \mathcal{F} is saturated if TX/\mathcal{F} has no torsion. Since \mathcal{F} is saturated, the Zariski closed subset $Sing(\mathcal{F}) \subset X$ over which \mathcal{F} is not a subbundle has codimension at least two, and any local section of \mathcal{F} defined on the complement of $Sing(\mathcal{F})$ extends across $Sing(\mathcal{F})$.

Definition 1.1. *We say that \mathcal{F} is a foliation if, for any two germs of sections V, W of \mathcal{F} , their Lie Bracket $[V, W]$ is also a germ of section of \mathcal{F} . Equivalently, the Lie bracket defines a morphism of sheaves of \mathcal{O}_X -modules ${}^1L : \wedge^2 \mathcal{F} \rightarrow TX/\mathcal{F}$ which vanishes identically on X .*

This is the ‘Frobenius’ integrability condition. It implies that, over $U_{\mathcal{F}} := X \setminus Sing(\mathcal{F})$, each point has a local analytic open neighbourhood $U = F \times B$, with F, B open subsets of \mathbb{C}^r and \mathbb{C}^{n-r} respectively such that $\mathcal{F}_U = Ker(d\beta)$, where $\beta : U \rightarrow B$ is the projection onto the second factor B . In particular, for each such $x = (f, b) \in U$, there is a unique germ of manifold $F_x = F \times \{b\}$ of dimension r , called the germ of the ‘leaf’ of \mathcal{F} at x , which has in each of its points $x' := (f', b)$, the subspace $\mathcal{F}_{x'}$ as its tangent space.

We thus define an equivalence relation on $U_{\mathcal{F}}$ for which two points are equivalent if they can be connected by a chain of germs of leaves of \mathcal{F} locally defined as above. The classes of this relation are called the leaves of \mathcal{F} . They are the maximal connected, r -dimensional manifolds immersed (but not necessarily closed) in $U_{\mathcal{F}}$, and tangent to \mathcal{F} . Near the points of $Sing(\mathcal{F})$, the leaves of \mathcal{F} may have a chaotic behaviour.

Example 1.2. *The simplest example of an everywhere regular foliation with non-closed leaves is the following: $X := \mathbb{C}^2/\Lambda$ is an Abelian surface, and $\mathcal{F} = X \times \{\mathbb{C}.v\} \subset TX = X \times \mathbb{C}^2$ is the trivial rank one subbundle generated by a vector $v \in \mathbb{C}^2$. Let $\Gamma := \mathbb{C}.v \cap \Lambda$: this is a closed additive subgroup of $\mathbb{C}.v$ of rank ρ either 0, 1 or 2. So all leaves are isomorphic to $\mathbb{C}.v/\Gamma$, which is isomorphic to either \mathbb{C}, \mathbb{C}^* , or an elliptic curve when $\rho = 2$, i.e when Γ is a cocompact sublattice of $\mathbb{C}.v$. If the leaves of \mathcal{F} are not compact, they are Zariski-dense, but their topological closures are real tori of real dimension d either 3 or 4. It may happen that $d = 3$ even if the leaves of \mathcal{F} are isomorphic to \mathbb{C} . This happens for example if $v = e_1 + \sqrt{2}.e_2$, $J.v = e_1 + \sqrt{3}.e_3 := w$, the lattice defining X being generated over \mathbb{Z} by e_1, \dots, e_4 , and the complex*

¹Indeed: $[fV, gW] = fg.[V, W] + (fV(g).W - g.W(f).V) \equiv fg.[V, W]$ modulo \mathcal{F} for any local sections V, W of \mathcal{F} , and holomorphic functions f, g .

structure J on \mathbb{R}^4 extending the one defined above ($J.v = w$). One obtains so a two-dimensional compact complex torus X . It is less obvious whether this can be done in such a way that X is an abelian variety.

Example 1.3. Let $X := \mathbb{P}^2$, with affine coordinates (x, y) and \mathcal{F} be defined as the Kernel of the (rational) 1-form $w := dy - \lambda dx$, where $\lambda \in \mathbb{C}^*$. The leaves are then defined by the equation $y = x^\lambda$, and are thus algebraic if and only if $\lambda \in \mathbb{Q}^*$. The Zariski and topological closures of the leaves can be easily described in terms of λ . Examples 1.2 and 1.3 show that algebraicity is an arithmetic property in families of foliations.

Example 1.4. The globally simplest foliations, however, are the ones arising from dominant rational fibrations $f : X \dashrightarrow Z$, where Z is a manifold of dimension $p := n - r$, and the generic fibre of f is connected. The associated foliation is then $\mathcal{F} := (\text{Ker}(df))^{\text{sat}}$, where \bullet^{sat} denotes the saturation inside TX , and $\text{Sing}(\mathcal{F})$ is the union of the indeterminacy locus of f , and of the singularities of the (reduction of the) fibres.

Definition 1.5. Let $\mathcal{F} \subset TX$ be a foliation on X . We say that \mathcal{F} is ‘algebraic’ if all leaves of \mathcal{F} are algebraic submanifolds of X , that is: if their Zariski and topological closures coincide, or equivalently, have the same dimension r .

Using Chow-scheme theory, one shows that \mathcal{F} is algebraic if and only if it arises from a fibration, as in Example 1.4.

Problem: Find ‘numerical’ criteria for the algebraicity of foliations $\mathcal{F} \subset TX$. The criteria we shall give and use below are the positivity of intersection numbers of $\det(\mathcal{F})$ and ‘movable’ classes of curves on either X or the projectivisation of \mathcal{F} . We shall see that these conditions also imply restrictions on the structure of the leaves of \mathcal{F} .

The criterion we shall prove and use here is the following, inspired by, and partially extending, former results ([28], [6], [11]):

Theorem 1.6. ([10], [15]) *The foliation $\mathcal{F} \subset TX$ is algebraic if the dual $\mathcal{F}^* := \text{Hom}_{\mathcal{O}_X}(\mathcal{F}, \mathcal{O}_X)$ of \mathcal{F} is not pseudo-effective².*

We also have the next particular case:

Theorem 1.7. *Let $\mathcal{F} \subset TX$ be a foliation with $\mu_{\alpha, \min}(\mathcal{F}) > 0$ for some $\alpha \in \text{Mov}(X)$. Then \mathcal{F}^* is not pseudo-effective, \mathcal{F} is an algebraic foliation. Moreover, its leaves have rationally connected closures.*

The notions of rational connectedness and $\mu_{\alpha, \min}$ will be defined in sections 4 and 6 below. We now define the pseudo-effectivity.

²This is proved, but not stated explicitly in [10]. It is stated explicitly in [15].

Definition 1.8. A coherent sheaf \mathcal{G} on X , is pseudo-effective if, for any $j > 0, c > 0$, $H^0(X, \text{Sym}^{[m]}(\mathcal{G}) \otimes A^j) \neq \{0\}$ for some $m > j.c$. Here $\text{Sym}^{[m]}(\mathcal{G})$ denotes the reflexive hull (i.e: the double dual) of $\text{Sym}^m(\mathcal{G})$.

Otherwise, if $H^0(X, \text{Sym}^{[m]}(\mathcal{G}) \otimes A^j) = \{0\}, \forall m > c(A).j$, any $j > 0$, and some constant $c(A) > 0$, \mathcal{G} is said to be not pseudo-effective.

Equivalently, by the next remark, \mathcal{G} is not pseudo-effective if \mathcal{G} is locally free on $U = X \setminus S$, where $S \subset X$ is Zariski closed of codimension at least 2, and if $H^0(U, \text{Sym}^m(\mathcal{G}) \otimes A^j) = \{0\}, \forall m > c(A).j$.

Remark 1.9. Recall that \mathcal{F} is reflexive if the natural morphism $\mathcal{F} \rightarrow \mathcal{F}^{**}$ is an equality. Such a sheaf is torsionfree and normal (i.e: any section defined on a big Zariski open subset $U \subset X$ extends to X). If $\text{rk}(\mathcal{F}) = 1$, reflexive means locally free on a smooth X . If $\mathcal{F} \subset \mathcal{E}$ is of rank one, with \mathcal{E} locally free, then \mathcal{F} is reflexive if and only if it is saturated in \mathcal{E} (but not necessarily a subbundle). Its powers $\mathcal{F}^k \subset \text{Sym}^k \mathcal{E}, k > 0$ are then all saturated.

We shall see in §3 and §4 how to check the non-pseudo-effectivity of a sheaf \mathcal{G} using negativity or non-positivity of intersection numbers with ‘movable classes of curves’.

The following consequences of Theorem 1.6 can be stated without reference to pseudo-effectivity.

Corollary 1.10. ([10],[11]) Let s be a nonzero section of $\otimes^m TX \otimes L$, for some $m > 0$ and L a line bundle with $c_1(L) = 0$. Assume that s vanishes somewhere on X . Then X is uniruled (i.e: covered by rational curves).

Recall that a line bundle on X is said to be ‘big’ if $h^0(X, m.L) \sim C.m^n$, for some $C = C(X, L) > 0$, and $m \rightarrow +\infty$. A line bundle L is big if and only if $mL = A + E$ for some $m > 0$, A ample and E effective.

Corollary 1.11. ([10]) Let L be big line bundle on X , and assume there exists a sheaf injection $L \subset \otimes^m \Omega_X^1$, for some $m > 0$. Then K_X is big.

Remark 1.12. The statement of Corollary 1.11 is a form of stability specific to vector bundles $E := \otimes^m \Omega_X^1$, and fails for general E : on any positive-dimensional X , if $E := A \oplus A^{\otimes -2}$, with $L = A$ ample of rank one, $\det(E) = -A$, although A injects in E .

Products $X = \mathbb{P}^1 \times Z_{n-1}$, with Z_{n-1} of general type, $f : X \rightarrow Z$ the projection, and $L = f^*(K_Z) \subset \Omega_X^{n-1}$, with $\kappa(X, L) = (n-1)$ submaximal, show that K_X may not be pseudo-effective. The positivity of subsheaves $L \subset \otimes^m \Omega_X^1$ is preserved by K_X only when L is big.

We shall formulate a more general version of ‘birational stability’ for tensor powers of cotangent bundles in §.7.

We shall apply in §8 the Corollary 1.11 to moduli of canonically polarized manifolds.

2. PROOF OF THE ALGEBRAICITY CRITERION.

Let $S := \text{Sing}\mathcal{F} \subset X$ be the Zariski closed set of codimension at least 2 over which \mathcal{F} is not a subbundle of TX , and $U_{\mathcal{F}} := X \setminus S$. For any $x \in U_{\mathcal{F}}$, let $U_x = F_x \times B_x$ be an open neighbourhood U_x of x in X such that the leaves of $\mathcal{F}|_{U_x}$ are the fibres of the projection $\beta_x : U_x \rightarrow B_x$. If $\Delta \subset X \times X$ is the diagonal of X , and if $\Delta_{\mathcal{F}}$ is the Zariski open set of Δ mapped to $U_{\mathcal{F}}$ by (either) projections $p_i : X \times X \rightarrow X, i = 1, 2$, then there exists a germ W of submanifold of dimension $n + r$ of $U_{\mathcal{F}} \times U_{\mathcal{F}}$ such that for each $x \in U_{\mathcal{F}}$, $W \cap U_x \times U_x = U_x \times_{B_x} U_x$. In other words, the fibre W_x over $x \in U_{\mathcal{F}}$ of the first projection $p_{1|W} : W \rightarrow U_{\mathcal{F}}$ is a germ of the leaf of \mathcal{F} through x .

- The leaves of \mathcal{F} will thus be algebraic if (and only if) the manifold W is a germ of an algebraic subvariety of $X \times X$, or equivalently, if the Zariski closure V of W has the same dimension $n + r$ as W . Let $d := \dim(V)$; if B is any ample line bundle on V , there exists a constant $C = C(B, V) > 0$ such that $h^0(V, k.B) \cong C.k^d$, when $k \rightarrow +\infty$. We thus only need to prove that $h^0(V, k.B) \leq C.k^{n+r}$ for some $C > 0$, and some ample B on V in order to prove Theorem 1.6.

- Notice next that it is sufficient to show that $h^0(W, k.B) \leq C.k^{n+r}$ for some $C > 0$, since the restriction map $\text{res} : H^0(V, k.B) \rightarrow H^0(W, k.B)$ is injective by the Zariski density of W in V . We now prove this last inequality.

- Let thus B be ample on $X \times X$. Any section s of $k.B$ on W admits a unique development in power series along the fibres of $p_{1|W} : W \rightarrow U_{\mathcal{F}}$ of the form $s = \sum_{m \geq 0} s_{m,k}$, where $s_{m,k} \in H^0(U_{\mathcal{F}}, \text{Sym}^m(\mathcal{F}^*) \otimes k.B), \forall m, k$.

Since \mathcal{F}^* is not pseudo-effective, $H^0(U_{\mathcal{F}}, \text{Sym}^m(\mathcal{F}^*) \otimes k.B) = \{0\}$ if $m > C'.k$, for some $C' = C'(B)$.

Thus $h^0(W, k.B) \leq \sum_{m=0}^{m=C'.k} h^0(U_{\mathcal{F}}, \text{Sym}^m(\mathcal{F}^*) \otimes k.B)$.

If $U_{\mathcal{F}} = X$ (i.e. \mathcal{F} is a subbundle of TX), $h^0(U_{\mathcal{F}}, \text{Sym}^m(\mathcal{F}^*) \otimes k.B) \leq C''.(m+k)^{n+r-1}, \forall k > 0, \forall m \geq 0$, and some $C'' > 0$. Indeed, $h(m, k) := h^0(X, \text{Sym}^m(\mathcal{F}^*) \otimes k.B) = h^0(\mathbb{P}(\mathcal{F}), m.L + k.B')$, where $L = \mathcal{O}_{\mathbb{P}(\mathcal{F})}(1)$, B' is the pullback of B , and so $h(m, k) \leq C''.(m+k)^{n+r-1}$, since $\dim(\mathbb{P}(\mathcal{F})) = n + r - 1$, by dominating L, B' by an ample line bundle A' on $\mathbb{P}(\mathcal{F})$ such that $A' - B'$ and $A' - L$ are effective. Thus:

$$\sum_{m=0}^{m=C'.k} h^0(U_{\mathcal{F}}, \text{Sym}^m(\mathcal{F}^*) \otimes k.B) \leq \sum_{m=0}^{m=C'.k} C''.(m+k)^{n+r-1} \leq (C'.k+1).(C''.(C'.k+k)^{n+r-1}) \leq (C'+1)^{n+r} C'' .k^{n+r},$$

which concludes the proof of Theorem 1.6 if $U = X$.

The general case $U_{\mathcal{F}} \neq X$ is obtained by applying [31], III.5.10(3), which shows that the same inequalities $h^0(U_{\mathcal{F}}, \text{Sym}^m(\mathcal{F}^*) \otimes k.B) \leq C''.(m+k)^{n+r-1}, \forall k > 0, \forall m \geq 0$, and some $C'' > 0$ still hold true, by constructing a suitable modification of $\mathbb{P}(\mathcal{F})$, and of $\mathcal{O}_{\mathbb{P}(\mathcal{F})}(1)$.

More precisely: if $p : P := \mathbb{P}(\mathcal{F}) := \text{Proj}(\text{Sym}^{\bullet} \mathcal{F}) \rightarrow X$, with $L := \mathcal{O}_P(1)$ the tautological line bundle on P , $p_*(L^m) = \text{Sym}^m(\mathcal{F}), \forall m > 0$.

Moreover, if $\rho : P' \rightarrow P$ is a resolution of the singularities of P which coincides with P over $U_{\mathcal{F}}$, there exists an effective, ρ -exceptional divisor $E \subset P'$ such that $(p \circ r)_*(m.(L + E)) = \text{Sym}^{[m]}(\mathcal{F}), \forall k > 0$. See Proposition 3.9 for a simplified proof.

3. PSEUDOEFFECTIVITY AND MOVABLE CLASSES.

Definition 3.1. *Let $(C_t)_{t \in T}$ be an algebraic family of curves parametrised by an irreducible projective variety T . Assume that C_t is irreducible for $t \in T$ generic, and that X is covered by the union of the C'_t s. Then $\alpha := [C_t] \in H^{2n-2}(X, \mathbb{Z})$ is independent of $t \in T$. We call such a class a **geometrically movable class** on X .*

*The closed convex cone of $H^{2n-2}(X, \mathbb{R})$ generated by the geometrically movable classes is called the **movable cone** of X , denoted $\text{Mov}(X)$, and its elements are the **movable classes** of X . From [7], we even have that $\alpha \in H^{n-1, n-1}(X, \mathbb{R})$ is in $\text{Mov}(X)$ if and only if $\alpha.D \geq 0$ for any irreducible effective divisor D on X .*

Example 3.2. *1. If A is ample on X , then $A^{n-1} \in \text{Mov}(X)$. There are many more examples, such as the following:*

2. If $C \subset X$ is an irreducible curve, locally complete intersection, with ideal sheaf I_C and ample normal bundle $(I_C/I_C^2)^$, then $\alpha := [C] \in \text{Mov}(X)$. Indeed: for any irreducible effective divisor D , we have $D.C \geq 0$, since $D|_C$ is a quotient of $(I_C/I_C^2)^*$. See [33], [35] for more on this situation.*

Remark 3.3. *If D is an effective \mathbb{Q} -divisor on X , $D.\alpha \geq 0, \forall \alpha \in \text{Mov}(X)$. More generally, if $m.D + A$ is effective for infinitely many $m > 0$, and some ample A , $D.\alpha \geq 0, \forall \alpha \in \text{Mov}(X)$. Said otherwise: if D is pseudo-effective, then $D.\alpha \geq 0, \forall \alpha \in \text{Mov}(X)$.*

An important result is the following converse.

Theorem 3.4. ([7],[31]) *Let L be a line bundle on X . Then L is pseudo-effective if and only if $L.\alpha \geq 0, \forall \alpha \in \text{Mov}(X)$.*

*Equivalently, L is **not** pseudo-effective if and only if $L.\alpha < 0$ for some $\alpha \in \text{Mov}(X)$.*

Pseudo-effectivity is fundamental in birational geometry when applied to $L = K_X$ because of the next:

Theorem 3.5. ([29]) *If K_X is not pseudo-effective, then X is uniruled (i.e: covered by rational curves).*

The only known proof rests on positive characteristic techniques. The original statement of Miyaoka-Mori is: if $(C_t)_{t \in T}$ is an algebraic family of curves (of possibly large genus) such that $K_X.C_t < 0$, it is possible to find such a family where the C'_t s are rational curves!

Remark 3.6. *There exists (Nagata, Mumford, see [30]) divisors D such that $D.\alpha \geq 0, \forall \alpha \in \text{Mov}(X)$, but which are not \mathbb{Q} -effective. The main conjecture of birational geometry is that K_X is \mathbb{Q} -effective if $K_X.\alpha \geq 0, \forall \alpha \in \text{Mov}(X)$.*

The following birational invariance is crucial, here.

Corollary 3.7. *Let $\pi : X' \rightarrow X$ be a birational morphism between smooth projective manifolds.*

1. *Let $\alpha \in \text{Mov}(X)$, then $\pi^*(\alpha) \in \text{Mov}(X')$.*
2. *Let $\alpha' \in \text{Mov}(X')$, then $\pi_*(\alpha') \in \text{Mov}(X)$.*
3. *$\alpha \in \text{Mov}(X)$ if and only if $\alpha' := \pi^*(\alpha) \in \text{Mov}(X')$.*

Proof. The first (resp. second) claim follows from the equality: $\pi^*(\alpha).D' = \alpha.\pi_*(D')$ for any divisor D' on X' (resp. $\pi^*(\alpha).\pi^*(D) = \pi^*(\alpha).(D' + E') = \pi^*(\alpha).D'$, where D' is the strict transform of D , and E' is π -exceptional, since $\pi^*(\alpha).E' = 0, \forall E'$, π -exceptional divisor on X'). The third claim follows from the others since $\pi_*(\pi^*(\alpha)) = \alpha, \forall \alpha$. \square

The following is an immediate consequence of Theorem 3.4 since $p^*(A) + \varepsilon.\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$ is ample on $\mathbb{P}(\mathcal{E})$ for any $\varepsilon > 0$ sufficiently small, and $\text{Sym}^m(\mathcal{E}) = p_*(\mathcal{O}_{\mathbb{P}(\mathcal{E})}(m)), \forall m > 0$:

Corollary 3.8. *Let \mathcal{E} be a rank r vector bundle on X . Then \mathcal{E} (identified with its sheaf of sections) is pseudo-effective if and only if $L := \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$ is pseudo-effective on $p : \mathbb{P}(\mathcal{E}) \rightarrow X$, that is, if $L.\alpha \geq 0, \forall \alpha \in \text{Mov}(\mathbb{P}(\mathcal{E}))$.*

The above Corollary 3.8 extends to the case when $\mathcal{G} = E$ is not locally free by considering a suitable modification of $\mathbb{P}(\mathcal{G})$ as constructed in [31, III,5.10]. We give a simplified proof, following the strategy of [31], in Proposition 3.9 below (to be applied to a smooth model P of the main component of $\mathbb{P}(\mathcal{F})$, as in [31, V,3.23]).

Proposition 3.9. *Let $p : P \rightarrow X$ be a fibration with X normal, P smooth, both projective, let D be a Cartier divisor on P such that $p_*(\mathcal{O}_P(D)) \neq 0$.*

1. *If E is an effective divisor on P supported on the exceptional divisor $\text{Exc}(p)$ of p , and such that, for every divisorial component Γ of $\text{Exc}(p)$, its restriction E_Γ to Γ is not pseudo-effective, then $p_*(\mathcal{O}_P(k.(D + \ell.E))) = p_*(\mathcal{O}_P(k.D))^{**}, \forall k > 0$, where $\ell > 0$ is chosen to be sufficiently large, so that $(D + \ell.E)_\Gamma$ is not pseudo-effective for any Γ .*
2. *There exists a divisor E satisfying the previous properties.*

Proof. 1. Let $\Sigma \subset X$ be the codimension 2 or more locus of points over which p is not equidimensional, so that $\text{Exc}(p) \subset p^{-1}(\Sigma)$. There is some $\ell > 0$ such that $(D + \ell.E)_\Gamma$ is not pseudo-effective for every Γ , since this is the case for E . We replace E by $\ell.E$, so that $\ell = 1$, to simplify notations. Then every section of $\mathcal{O}_P(k.(D + E))$ vanishes on

$Exc(p)$, for every $k > 0$, and has thus no pole there. Every element s of $H^0(P, p^*(p_*(\mathcal{O}_P(k.D))^{**}))$ lies in $H^0(P, k.(D + E) + E')$, for some divisor E' supported on $Exc(p)$, which may be supposed to be effective. But then $E' \leq t.E$ for some $t > 0$ since the support of E coincides with the divisorial part of $Exc(p)$ by our non pseudo-effectivity assumption. Since $t.E$ is not pseudo-effective on each Γ , $H^0(P, k.(D + E) + t.E) = H^0(P, k.(D + E)) = H^0(P, p^*(p_*(\mathcal{O}_P(k.D))^{**}))$, which implies the claim.

2. Let $v : X' \rightarrow X$ be a modification, $q : P' \rightarrow X'$, $u : P' \rightarrow P$ be the normalisation of the main component of $P \times_X X'$, so that $p \circ u = v \circ q$. We choose v to be projective birational, with X' smooth, and moreover such that q is equidimensional, and such that there exists an effective divisor $\Delta \subset Exc(v)$ such that $-\Delta$ is ample on the divisorial part of $Exc(v)$.

Let $E := u_*(q^*(\Delta))$, and let H be a sufficiently ample divisor on P , with H_Γ its restriction to Γ , for each divisorial component Γ of $Exc(p)$. Let $\alpha_\Gamma := H_\Gamma^{d-2}$, where $d := \dim(P)$, then $\alpha_\Gamma \in Mov(\Gamma)$. If $\Gamma' \subset P'$ is the strict transform of Γ by u , then $q^*(\Delta).u^*(\alpha_\Gamma) = E.\alpha_\Gamma$ since the generic member of the family α_Γ does not meet the indeterminacy locus of $u|_{\Gamma'} : \Gamma' \rightarrow \Gamma$. We thus just have to show that $q^*(\Delta).u^*(\alpha_\Gamma) < 0$.

Moreover, $q^*(\Delta).u^*(\alpha_\Gamma) = \Delta.q_*(u^*(\alpha_\Gamma)) < 0$. Indeed: $q_*(u^*(\alpha_\Gamma)) \in Mov(q(\Gamma'))$, $-\Delta$ is ample on the divisorial part of $Exc(v)$, and $q(\Gamma')$ is a divisorial component of $Exc(v)$, by the equidimensionality of q . Thus $E.\alpha_\Gamma < 0$, and E_Γ is not pseudo-effective. \square

The description of $Mov(\mathbb{P}(E))$ is, in general, quite delicate. An important particular situation where the pseudo-effectivity can be tested on X rather than on $\mathbb{P}(E)$ is exposed in the next section.

4. HARDER-NARASIMHAN FILTRATIONS AND PSEUDO-EFFECTIVITY.

The data here are $X, \mathcal{G}, \alpha \in Mov(X), r := rk(\mathcal{G}) > 0$ as above. We assume always \mathcal{G} to be nonzero, torsionfree. See [25], Chap. V and [21] for a detailed treatment.

The (always locally free, since X is smooth), rank-one, sheaf $det(\mathcal{G})$ is defined as $det(\mathcal{G}) := \wedge^r(\mathcal{G})^{**}$.

Definition 4.1. *The slope of \mathcal{G} with respect to α is: $\mu_\alpha(\mathcal{G}) := \frac{det(\mathcal{G}).\alpha}{r}$. We say that \mathcal{G} is α -stable (resp. α -semi-stable) if: $\mu_\alpha(\mathcal{H}) < \mu_\alpha(\mathcal{G})$ (resp. $\mu_\alpha(\mathcal{H}) \leq \mu_\alpha(\mathcal{G})$), $\forall \mathcal{H} \subsetneq \mathcal{G}$ coherent subsheaf.*

Notice that if $\mathcal{F} \subsetneq \mathcal{G}$ is a saturated subsheaf of rank r with torsionfree quotient \mathcal{Q} of rank s , we have: $det(\mathcal{G}) = det(\mathcal{F}) + det(\mathcal{Q})$ and so:

$$(*) \mu_\alpha(\mathcal{G}) = \frac{r}{r+s}.\mu_\alpha(\mathcal{F}) + \frac{s}{r+s}.\mu_\alpha(\mathcal{Q}) \in [\mu_\alpha(\mathcal{F}), \mu_\alpha(\mathcal{Q})].$$

If \mathcal{F} is torsionfree, then $\mu_\alpha(\mathcal{F}) = \mu_\alpha(\mathcal{F}^{**}) = -\mu_\alpha(\mathcal{F}^*)$, since these equalities need to be checked only on a big Zariski-open subset of X .

These notions are classical when $\alpha = [A]^{n-1}$, where A is an ample line bundle. Just as in this classical case, we still have (with essentially the same proof, by induction on the rank, and the equality (*)):

Lemma 4.2. *For \mathcal{G}, α as above, there is a unique maximum $\mathcal{H} \subset \mathcal{G}$ with $\mu_\alpha(\mathcal{H})$ maximum (i.e: for any $\mathcal{H}' \subset \mathcal{G}$, we have: $\mu_\alpha(\mathcal{H}') \leq \mu_\alpha(\mathcal{H}) := \mu_{\alpha, \max}(\mathcal{G})$, in case of equality: $\mathcal{H}' \subset \mathcal{H}$).*

\mathcal{H} is the α -maximal destabilising subsheaf of \mathcal{G} , denoted $\mathcal{G}^{\alpha, \max}$. It is, by construction, α -semi-stable, and saturated inside \mathcal{G} , so its quotient $\mathcal{G}_{\alpha, \min} := \mathcal{G}/\mathcal{G}^{\alpha, \max}$ is torsionfree (or zero iff \mathcal{G} is semi-stable).

Proof. Obvious if $\text{rank}(\mathcal{G}) = 1$, and so for direct sum of copies of a line bundle. Then embed \mathcal{G} in the direct sum of a certain number of copies of A , sufficiently ample. Then $\mu_\alpha(\mathcal{G}') \leq \mu_\alpha(\oplus^N A) = \mu_\alpha(A)$ for any $\mathcal{G}' \subset \mathcal{G}$. This proves the boundedness of $\mu_\alpha(\mathcal{G}')$, for $\mathcal{G}' \subset \mathcal{G}$. If the class α is rational, one easily gets the existence of a maximum for these slopes, because of the finiteness of the possible denominators, and one can choose the rank to be maximal. The conclusion then follows. If the class α is not rational, one needs a further (but still elementary) argument: choose $\mathcal{G}_i, i = 1, 2$ both different from their intersection, of the same maximal rank r with α -slopes $\mu_i, i = 1, 2$, approaching from below up to $\varepsilon > 0$ the upper bound $\mu := \mu_{\alpha, \max}(\mathcal{G})$ of α -slopes of all $\mathcal{G}' \subset \mathcal{G}$. The sum $\mathcal{G}_1 + \mathcal{G}_2$ has rank larger than r and slope at least each $\mu - 2r\varepsilon$ (by a simple computation using the exact sequence $0 \rightarrow \mathcal{G}_1 \cap \mathcal{G}_2 \rightarrow \mathcal{G}_1 \oplus \mathcal{G}_2 \rightarrow \mathcal{G}_1 + \mathcal{G}_2 \rightarrow 0$). Contradiction to the maximality of r for the ranks of $\mathcal{G}' \subset \mathcal{G}$ with slope at least $\mu - 2r\varepsilon$ if $\varepsilon > 0$ is chosen sufficiently small. \square

Corollary 4.3. *There are a unique integer $s \geq 0$, and an increasing filtration $\{0\} = \mathcal{H}_0 \subsetneq \mathcal{H}_1 \subsetneq \mathcal{H}_2 \subsetneq \dots \subsetneq \mathcal{H}_s = \mathcal{G}$ by saturated subsheaves such that: $\mathcal{H}_{j+1}/\mathcal{H}_j$ is α -semistable for $j = 0, \dots, s-1$, and $\mu_\alpha(\mathcal{H}_{j+1}/\mathcal{H}_j) > \mu_\alpha(\mathcal{H}_{j+2}/\mathcal{H}_{j+1})$ for $j = 0, \dots, s-2$.*

This filtration is called the α -Harder-Narasimhan filtration of \mathcal{G} . We write $\mu_{\alpha, \min}(\mathcal{G}) := \mu_\alpha(\mathcal{G}_{\alpha, \min}) := \mu_\alpha(\mathcal{G}/\mathcal{H}_{s-1})$, and $\mu_{\alpha, \max}(\mathcal{G}) := \mu_\alpha(\mathcal{G}^{\alpha, \max})$.

Proof. Induction on s applied to $\mathcal{G}/\mathcal{G}^{\alpha, \max}$ and $\mathcal{H}_1 = \mathcal{G}^{\alpha, \max}$. \square

Lemma 4.4. *1. Let $\mathcal{H} \neq \{0\}$ be a torsionfree quotient sheaf of \mathcal{G} on X . Then: $\mu_\alpha(\mathcal{H}) \geq \mu_{\alpha, \min}(\mathcal{G})$. If equality holds, \mathcal{H} is a quotient of \mathcal{G}_{\min} .*

2. Let \mathcal{F}, \mathcal{G} be torsionfree coherent sheaves on X . Assume that $\mu_{\alpha, \min}(\mathcal{F}) > \mu_{\alpha, \max}(\mathcal{G})$. Then $\text{Hom}(\mathcal{F}, \mathcal{G}) = \{0\}$.

3. If $\det(\mathcal{G}) \cdot \alpha > 0$, then $\mu_{\alpha, \max}(\mathcal{G}) = \mu_{\alpha, \min}(\mathcal{G}^{\alpha, \max}) > 0$.

Proof. 1. Induction on r : if \mathcal{H} is a nonzero quotient of \mathcal{G} , it fits in a short exact sequence $0 \rightarrow K \rightarrow \mathcal{H} \rightarrow \mathcal{Q} \rightarrow 0$, in which K (resp. \mathcal{Q}) is a quotient of $\mathcal{G}^{\alpha, \max}$ (resp. $\mathcal{G}/\mathcal{G}^{\alpha, \max}$), hence the conclusion since

both K and \mathcal{Q} have slope at least $\mu_{\alpha, \min}(\mathcal{G})$. The second claim holds by induction on s , because $\mu_{\alpha, \min}(\mathcal{G}/\mathcal{G}^{\alpha, \max}) > \mu_{\alpha, \min}(\mathcal{G})$ if \mathcal{G} is not semi-stable.

2. If $0 \neq h \in \text{Hom}(\mathcal{F}, \mathcal{G})$, its image \mathcal{H} a subsheaf of \mathcal{G} , and a quotient of \mathcal{F} . We thus get (with $\mu = \mu_{\alpha}$): $\mu_{\max}(\mathcal{G}) \geq \mu(\mathcal{H}) \geq \mu_{\min}(\mathcal{F})$, a contradiction.

3. is obvious. \square

Lemma 4.5. *We have: $(\mathcal{G}^*)^{\alpha, \max} = (\mathcal{G}_{\alpha, \min})^*$, in particular: $\mu_{\alpha, \min}(\mathcal{G}) = -\mu_{\alpha, \max}(\mathcal{G}^*)$, and $\text{HN}_{\alpha}(\mathcal{G}^*) = (\text{HN}_{\alpha}(\mathcal{G}))^*$ (i.e.: the terms for \mathcal{G}^* are the dual of those for \mathcal{G} in reverse order).*

Proof. Dualising the projection $\mathcal{G} \rightarrow \mathcal{G}_{\min}$, we get: $(\mathcal{G}_{\min})^* \subset \mathcal{G}^*$, and so $-\mu_{\min}(\mathcal{G}) \leq \mu_{\max}(\mathcal{G}^*)$. Dualising $(\mathcal{G}^*)^{\alpha, \max} \subset \mathcal{G}^*$, we get a generically surjective morphism $\mathcal{G} \rightarrow \mathcal{G}^{**} \rightarrow ((\mathcal{G}^*)^{\alpha, \max})^*$ from which the inequality $-\mu_{\max}(\mathcal{G}^*) \geq \mu_{\min}(\mathcal{G})$ follows. We thus get the equality $-\mu_{\max}(\mathcal{G}^*) = \mu_{\min}(\mathcal{G})$. Hence the second claim. The first claim follows from the second part of Lemma 4.4.(1). The last claim is seen by induction on the number of terms of the HN filtration. \square

Let $\mathcal{F} \hat{\otimes} \mathcal{G} := (\mathcal{F} \otimes \mathcal{G})^{**}$ be the reflexive tensor product of \mathcal{F} and \mathcal{G} .

Theorem 4.6. *For $X, \mathcal{G}, \mathcal{H}, \alpha$ as before, we have:*

1. $\mu_{\alpha, \min}(\mathcal{G}) = -\mu_{\alpha, \max}(\mathcal{G}^*)$, $(\mathcal{G}_{\alpha, \min})^* = (\mathcal{G}^*)^{\alpha, \max}$.
2. $(\mathcal{G} \hat{\otimes} \mathcal{H})^{\alpha, \max} = \mathcal{G}^{\alpha, \max} \hat{\otimes} \mathcal{H}^{\alpha, \max}$.
3. $\mu_{\alpha, \max}(\mathcal{G} \hat{\otimes} \mathcal{H}) = \mu_{\alpha, \max}(\mathcal{G}) + \mu_{\alpha, \max}(\mathcal{H})$.
4. $\mu_{\alpha, \max}(\hat{\otimes}^m(\mathcal{G})) = \mu_{\alpha, \max}(\widehat{\text{Sym}}^m(\mathcal{G})) = m \cdot \mu_{\alpha, \max}(\mathcal{G})$.
5. $\mu_{\alpha, \max}(\hat{\wedge}^p(\mathcal{G})) = p \cdot \mu_{\alpha, \max}(\mathcal{G})$, $\forall p > 0$.

The first claim follows from the fact that dualisation exchanges sub-objects and quotients. By contrast, the proof of claim 3 is quite deep, relying on the Kobayashi-Hitchin correspondance between α -stable bundles and Hermite-Einstein vector bundles (with respect to Gauduchon metrics). For this correspondance, originally due to Li-Yau in [26], see details in the book of Lübke-Teleman [27].

Lemma 4.7. *Let \mathcal{F} be a torsionfree coherent sheaf on X , and $\alpha \in \text{Mov}(X)$ such that $\mu_{\alpha, \min}(\mathcal{F}) > 0$. Then \mathcal{F}^* is not pseudo-effective.*

Proof. Let A be ample on X . Applying Theorem 4.6, we see that $-\mu := \mu_{\alpha, \max}(\mathcal{F}^*) < 0$, and so $\mu_{\alpha, \max}(\text{Sym}^{[m]}(\mathcal{F}) \otimes A^j) = m \cdot \mu + j \cdot A \cdot \alpha < 0$ for $m > j \cdot c$, where $c := \frac{A \cdot \alpha}{\mu}$, which implies that $h^0(X, \text{Sym}^{[m]}(\mathcal{F}) \otimes A^j) = 0$, as claimed. \square

The following criterion for foliations among distributions is crucial here:

Corollary 4.8. ([28]) *Let $\mathcal{F} \subset TX$ be a coherent subsheaf. If, for some $\alpha \in \text{Mov}(X)$, $2 \cdot \mu_{\alpha, \min}(\mathcal{F}) > \mu_{\alpha, \max}(TX/\mathcal{F})$, then \mathcal{F} is a foliation.*

Proof. Let $\Lambda : \wedge^2(\mathcal{F}) \rightarrow TX/\mathcal{F}$ be the sheaf morphism induced by the Lie bracket. The slopes assumption implies by claim 5 of Theorem 4.6 and claim 2 of Lemma 4.4 that it vanishes. The conclusion follows by Frobenius theorem. \square

From Theorem 1.6, and Lemmas 4.7 and 4.8, we get:

Theorem 4.9. *Let $\mathcal{F} \subset TX$ be a distribution with $\mu_{\alpha, \min}(\mathcal{F}) > 0$ for some $\alpha \in \text{Mov}(X)$. Assume that $2 \cdot \mu_{\alpha, \min}(\mathcal{F}) \mu_{\alpha, \max}(TX/\mathcal{F})$ (this is satisfied if \mathcal{F} is a piece of the Harder-Narasimhan filtration of TX relative to α). Then \mathcal{F} is an algebraic foliation.*

Remark 4.10. *In the situation of Theorem 4.9, we shall prove in the next two sections that the leaves of \mathcal{F} have rationally connected closures.*

Corollary 4.11. *Assume that K_X is not pseudo-effective. There then exists on X a nonzero foliation $\mathcal{F} \subset TX$ such that $\mu_{\alpha, \min}(\mathcal{F}) > 0$, the dual \mathcal{F}^* is not pseudo-effective, and \mathcal{F} is algebraic.*

Proof. Just apply Theorem 4.9, choosing $\mathcal{F} := TX^{\alpha, \max}$. The last claim requires new notions and techniques presented in the next two sections. \square

Remark 4.12. *We thus obtain the existence of (all of the) fibrations with rationally connected fibres on any given X with K_X not pseudo-effective without using the Minimal Model Program.*

Another easy, but central, property is the birational invariance ‘up’ of the condition $\mu_{\alpha, \min}(\mathcal{F}) > 0$ for subsheaves of the tangent bundles:

Lemma 4.13. *Let $\mathcal{F} \subset TX$ be a subsheaf, let $\pi : X' \rightarrow X$ be a birational morphism between X, X' projective smooth. Let $\mathcal{F}' \subset TX'$ be the pullback distribution, defined as: $\mathcal{F}' := \pi^*(\mathcal{F}) \cap TX'$. Let $\alpha \in \text{Mov}(X), \alpha' := \pi^*(\alpha)$. Then $\mu_{\alpha'}(\mathcal{F}') = \mu_{\alpha}(\mathcal{F})$, and $\mu_{\alpha', \min}(\mathcal{F}') = \mu_{\alpha, \min}(\mathcal{F})$.*

Proof. The statements hold for $\pi^*(\mathcal{F})$, hence for \mathcal{F}' since $\det(\mathcal{F}')$ and $\det(\pi^*(\mathcal{F}))$ coincide outside the exceptional divisor of π , and $\alpha' \cdot E = 0$ for each component E of this divisor. \square

Remark 4.14. *The statement does not (always) hold ‘down’ (i.e. if $\alpha = \pi_*(\alpha')$, but $\alpha' \neq \pi^*(\alpha)$): let $X = \mathbb{P}^2$, let \mathcal{F} be tangent to the conics C going through 4 points in general position, and let α' be the class of the strict transforms C' of these conics on the blow-up X' of \mathbb{P}^2 in the 4 given points. Then $\mu_{\alpha'}(\mathcal{F}') = 2 > 0$, since $\mathcal{F}'_{|C'} = -K_{C'}$, but $\mu_{\alpha}(\mathcal{F}) = 2 - 4 = -2 > 0$.*

5. PSEUDO-EFFECTIVITY OF RELATIVE CANONICAL BUNDLES.

Let $f : X \rightarrow Z$ be a fibration between complex projective manifolds. Let X_z be its generic (smooth) fibre. Let $f^*(\Omega_Z^1)^{\text{sat}} \subset \Omega_X^1$ be the

saturation of $f^*(\Omega_Z^1)$, and denote by $f^*(K_Z)^+ := \det(f^*(K_Z)^{sat})$. Let finally $K_{X/Z}^- := K_X - f^*(K_Z)^+$.

If $\mathcal{F} := \text{Ker}((df)^{sat}) \subset TX$, we thus have: $\det(\mathcal{F}^*) = K_{X/Z}^-$, by the consideration of the determinants in the exact sequence: $0 \rightarrow (f^*(\Omega_Z^1))^{sat} \rightarrow \Omega_X^1 \rightarrow \mathcal{F}^* \rightarrow 0$, dual to: $0 \rightarrow \mathcal{F} \rightarrow TX \rightarrow TX/\mathcal{F} \rightarrow 0$.

Definition 5.1. *We say that f is ‘neat’ if:*

1. *The discriminant locus $D(f) \subset Z$ of singular fibres of f is a divisor of strict normal crossings.*
2. *$f^{-1}(D(f)) \subset X$ is also a divisor of simple normal crossings.*
3. *There exists an equidimensional fibration $f_0 : X_0 \rightarrow Z$ and a birational morphism $u : X \rightarrow X_0$ such that $f_0 \circ u = f$.*

By suitable blow-ups of Z and X , any fibration can be made ‘neat’, as a consequence of Hironaka’s resolution of singularities, and Hironaka’s or Raynaud’s flattening theorems.

Theorem 5.2. *Let $f : X \rightarrow Z$ be a fibration between complex projective manifolds. Let X_z be its generic (smooth) fibre. If f is ‘neat’, and if K_{X_z} is pseudo-effective, $K_{X/Z}^-$ is pseudo-effective.*

This result strengthens a weak version of Viehweg’s theorem on the weak positivity of direct images of pluricanonical sheaves. A particular case was also obtained by Miyaoka using deformations of rational curves, Mori’s bend and break, and positive characteristic methods.

Corollary 5.3. *Assume that K_X is not pseudo-effective, let $\alpha \in \text{Mov}(X)$ be such that $K_X \cdot \alpha < 0$. Let $\mathcal{F} := TX^{\alpha, max} \subset TX$, thus $\mu_{\alpha, min}(\mathcal{F}) > 0$, and so \mathcal{F} is a foliation, by corollary 4.11, and algebraic, by Theorem 1.6. Moreover, after suitable blow-ups of X , we have: if $f : X \rightarrow Z$ is a fibration such that $\mathcal{F} = \text{Ker}(df)^{sat} \subset TX$, then:*

1. *K_{X_z} is not pseudo-effective. More generally:*
2. *If $f = h \circ g$ for rational fibrations $g : X \rightarrow Y$ and $h : Y \rightarrow Z$, then K_{Y_z} is not pseudo-effective, Y_z being the generic fibre of h (on suitable birational neat models of g, h).*

Proof. We may blow-up X, Z_0 for any rational fibration $f_0 : X \dashrightarrow Z_0$ with generic fibre the closure of a generic leaf of \mathcal{F} in such a way that the birational model $f : X \rightarrow Z$ (we keep the notation X for the blown-up manifold, so as to save notations) so obtained is regular and ‘neat’. By Lemma 4.13, the property $\mu_{\alpha, min}(\mathcal{F})$ is preserved on the blown-up manifold by lifting both the foliation and the movable class in the natural manner, so that the Theorem 5.2 can be applied to the new $f : X \rightarrow Z$.

Claim 1. By contradiction, assume not. Then $K_{X/Z}^- = \det(\mathcal{F}^*)$ is pseudo-effective, by Theorem 5.2. Thus $0 \leq K_{X/Z}^- \cdot \alpha = \det(\mathcal{F}^*) \cdot \alpha = -\mu_{\alpha}(\mathcal{F}) < 0$. Contradiction.

Claim 2. By contradiction again, assume that K_{Y_z} is pseudo-effective. By applying Theorem 5.2 to h , $\det(\mathcal{H}^*)$, and so also $g^*(\det(\mathcal{H})^*)$ were pseudo-effective, with $\mathcal{H} := \text{Ker}(dh)^{\text{sat}} \subset TY$. Thus $\mu_\alpha(g^*(\mathcal{H}^*)) = -\mu_\alpha(g^*(\mathcal{H})) \geq 0$. On the other hand, the fibration $g|_{X_z} := g_z : X_z \rightarrow Y_z$ implies that $g^*(\mathcal{H})$ is a quotient of \mathcal{F} , so $\mu_\alpha(g^*(\mathcal{H})) \geq \mu_{\alpha, \min}(\mathcal{F}) > 0$. Contradiction. \square

Remark 5.4. *With (much) more work, it is possible to show more: for general $z \in Z$, there exists $\alpha_z \in \text{Mov}(X_z)$ such that $\mu_{\alpha_z, \min}(TX_z) > 0$. We shall prove this less directly, but more simply, by using the theory of rational curves in the next section.*

6. RATIONAL CURVES AND NON-PSEUDOEFFECTIVITY OF THE CANONICAL/COTANGENT BUNDLES.

Definition 6.1. *X is said to be uniruled (resp. rationally connected) if through any generic point (resp. any two generic points) of X pass an irreducible rational curve (i.e: a non-constant image of \mathbb{P}^1).*

Example 6.2. 1. \mathbb{P}^n is rationally connected, so also any X birational to \mathbb{P}^n , or rationally dominated by \mathbb{P}^n .

2. X is rationally connected if $-K_X$ is ample (i.e: if X is Fano).

3. Smooth hypersurfaces of \mathbb{P}^{n+1} of degree at most $n+1$ are Fano (by adjunction), hence rationally connected.

4. If $X \rightarrow Z$ is a fibration with Z and X_z rationally connected, then X is rationally connected (a deep result of [19]).

The present birational theory of rational curves on projective manifolds is based on [19] and the following two results:

Theorem 6.3. ([29]) *X is uniruled if (and only if) K_X is not pseudo-effective.*

Theorem 6.4. *There is a unique fibration $r : X \rightarrow R$ (called the Maximally Rationally Connected (or MRC-) fibration of X) such that:*

1. *its fibres are rationally connected.*

2. *its base is not uniruled (i.e: K_R is pseudo-effective).*

Although in general, $r : X \rightarrow R$ is rational (almost holomorphic, in the sense that its generic fibre does not meet the indeterminacy locus), since all the properties considered here are of birational nature, we may and shall assume that r is regular.

The two extreme cases are: $X = R$, or R is a point, meaning respectively that X is not uniruled, and that X is rationally connected.

Using Chow-scheme theory, one gets a relative version as well, on suitable birational models:

Corollary 6.5. *If $f : X \rightarrow Z$ is a fibration, there exists $g : X \rightarrow Y$ and $h : Y \rightarrow Z$ such that for general $z \in Z$, $g_z := g|_{X_z} : X_z \rightarrow Y_z$ is the MRC of X_z , and so K_{Y_z} is pseudo-effective.*

Theorem 6.6. *Let $\mathcal{F} \subset TX$ be a foliation such that $\mu_{\alpha, \min}(\mathcal{F}) > 0$, for some $\alpha \in \text{Mov}(X)$. Then \mathcal{F} is algebraic, and its leaves have rationally connected closures.*

Proof. The algebraicity of \mathcal{F} follows from Theorem 4.9. Let $f = h \circ g$ be the relative MRC of f , a suitable fibration such that $\mathcal{F} = \text{Ker}(df)^{\text{sat}}$. By Corollary 5.3.1, and Theorem 6.3, X_z is uniruled and so $\dim(Y) < \dim(X)$. By Corollary 5.3.2, K_{Y_z} is not pseudo-effective if $\dim(Y) > \dim(Z)$, contradicting Corollary 6.5. Thus $Y = Z$, and the claim. \square

Corollary 6.7. *The following are equivalent:*

1. X is uniruled.
2. K_X is not pseudo-effective.
3. $h^0(X, mK_X + A) = 0$ for any ample A and large m .
4. $\mu_{\alpha, \min}(\Omega_X^1) < 0$, for some $\alpha \in \text{Mov}(X)$.

We have: $1 \implies 2, 3 \implies 4 \implies 1$ (the last two implications by Corollaries 4.11 and 6.6 respectively).

Corollary 6.8. *The following are equivalent:*

1. X is rationally connected
2. Ω_X^p is not pseudo-effective for any $p > 0$.
3. $h^0(X, \text{Sym}^m(\Omega_X^1) \otimes A) = 0$ for any $p > 0$, A ample and m large.
4. $\mu_{\alpha, \max}(\Omega_X^1) < 0$ for some $\alpha \in \text{Mov}(X)$.

Proof. The implications $1 \implies 2, 3$ are easy by taking a rational curve with ample normal bundle in X , $3 \implies 4$ is seen by contradiction, from Theorem 6.4, taking $L := r^*(K_R) \subset \Omega_X^p$ if $p := \dim(R) > 0$ is a pseudo-effective line bundle, thus such that $0 < h^0(X, m.L + A) \leq h^0(X, \text{Sym}^m(\Omega_X^p) \otimes r^*(A))$ for infinitely many m 's. That $4 \implies 1$ follows from Theorem 6.6 applied to $\mathcal{F} = TX$. \square

Remark 6.9. 1. *Theorem 6.3 (resp. Corollary 6.8) claims that X is uniruled (resp. rationally connected) if and only if there is a covering family of curves (possibly of large genus) on which $\det(TX)$ (resp. TX) is ample.*

2. *The ‘uniruledness conjecture’ claims that X is uniruled if K_X is not \mathbb{Q} -effective. It implies that in the above two corollaries, the ample A could be removed in the assertions 3, and that ‘pseudo-effective’ could be replaced by ‘ \mathbb{Q} -effective’ in assertions 2.*

Corollary 6.10. ([10],[11]) *Let s be a nonzero section of $\otimes^m TX \otimes L$, for some $m > 0$ and L a line bundle with $c_1(L) = 0$. Assume that s vanishes somewhere on X . Then X is uniruled (i.e: covered by rational curves).*

Proof. It is sufficient to show that $\mu_{\alpha, \max}(TX) > 0$, or equivalently, that $\mu_{\alpha, \max}(\otimes^m TX \otimes L) > 0$ for some $\alpha \in \text{Mov}(X)$. Let L be the subsheaf of rank one of $\otimes^m TX \otimes L$ generated by s , and let α be the geometrically movable class defined by a family of ample curves going through

some point where s vanishes. Thus $0 < \mu_\alpha(L) \leq \mu_{\alpha, \max}(\otimes^m TX \otimes L)$ as claimed. \square

Corollary 6.11. *Let $C \subset X$ be an irreducible projective curve such that $TX|_C$ is ample. Then X is rationally connected.*

This was observed in [18] by a different approach.

Proof. Apply Theorem 6.6, observing that $[C] \in \text{Mov}(X)$, by considering the graph C'' of the embedding $j : C' \subset X$ in $C' \times X$, C' the normalisation of C , which has normal bundle $TX|_{C'}$, so that Corollary 6.8 applies, and $[C''] \in \text{Mov}(C' \times X)$, so that $[C] = p_*([C'']) \in \text{Mov}(X)$. \square

7. BIRATIONAL STABILITY OF THE COTANGENT BUNDLE

Theorem 7.1. ([10]) *Assume that K_X is pseudo-effective. Let Q be a torsionfree quotient of $\otimes^m \Omega_X^1$, for some $m > 0$. Then $\det(Q)$ is pseudo-effective.*

Proof. K_X being pseudo-effective, $\mu_{\alpha, \min}(\Omega_X^1) \geq 0, \forall \alpha \in \text{Mov}(X)$, by Corollary 6.7. By Theorem 4.6, $\mu_{\alpha, \min}(\otimes^m \Omega_X^1) = m \cdot \mu_{\alpha, \min}(\Omega_X^1) \geq 0, \forall \alpha \in \text{Mov}(X)$, which means that any torsionfree quotient of $\otimes^m(\Omega_X^1)$ is pseudo-effective. \square

Theorem 7.2. ([10]) *Let L a pseudo-effective line bundle on X . If there exists a nonzero sheaf morphism $\lambda : L \rightarrow \otimes^m \Omega_X^1 \otimes K_X^{\otimes p}$ for some $m \geq 0, p > 0$, K_X is pseudo-effective.*

Proof. We proceed by induction on $n = \dim(X)$, the case $n = 1$ being obvious. Assume K_X is not pseudo-effective, there then exists $\alpha \in \text{Mov}(X)$ such that $K_X \cdot \alpha < 0$, and $\mathcal{F} \subset TX$ a foliation such that $\mu_{\alpha, \min}(\mathcal{F}) > 0$. Thus \mathcal{F} is algebraic, of the form $\text{Ker}(df)^{\text{sat}}$ for some fibration $f : X \rightarrow Z$ with rationally connected fibres, by Theorem 6.6.

If $\dim(Z) = 0$, we get $\mathcal{F} = TX$, and so: $\mu_{\alpha, \min}(TX) > 0$, which implies that $0 \leq L \cdot \alpha \leq \mu_{\alpha, \max}((\otimes^m \Omega_X^1) \otimes K_X^{\otimes p}) = -m \cdot \mu_{\alpha, \min}(TX) + p \cdot K_X \cdot \alpha < 0$. Contradiction.

If $\dim(Z) > 0$, then $\dim(X_z) < n$ and we may apply induction. Let $\lambda_z := \lambda|_{X_z} : L_z \rightarrow (\otimes^m \Omega_X^1 \otimes K_X^{\otimes p})|_{X_z}$ be the restriction. The exact sequence on X_z : $0 \rightarrow T \rightarrow \Omega_{X|X_z}^1 \rightarrow \Omega_{X_z}^1 \rightarrow 0$, with T a trivial bundle of rank $\dim(Z)$, induces a filtration on $(\otimes^m \Omega_X^1 \otimes K_X^{\otimes p})|_{X_z}$ with graded pieces isomorphic to $T^{\otimes(m-j)} \otimes^j \Omega_{X_z}^1 \otimes K_{X_z}^{\otimes p}$, and we thus get for some $j \in \{0, \dots, m\}$ a nonzero sheaf morphism $L_{X_z} \rightarrow \otimes^j \Omega_{X_z}^1 \otimes K_{X_z}^{\otimes p}$. The induction then implies that K_{X_z} is pseudo-effective, since L_{X_z} is, like L on X , pseudo-effective on X_z , for z general in Z (i.e. outside countably many strict Zariski-closed subsets). This contradicts the non-pseudo-effectivity of K_X . \square

Corollary 7.3. ([10]) *Let L a big line bundle on X . If there exists a nonzero sheaf morphism $L \rightarrow \otimes^m \Omega_X^1$ for some $m > 0$, K_X is big.*

Proof. We may assume that L is saturated, let $Q := (\otimes^m \Omega_X^1/L)$. If K_X is pseudo-effective, $\det(\otimes^m \Omega_X^1) = N.K_X = L + \det(Q)$. Since L is big and $\det(Q)$ is pseudo-effective by Theorem 7.1, $N.K_X$, and so K_X , is big.

We show that K_X is pseudo-effective, which will imply the claim. Since L is big, $k.L + K_X$ is effective for some large $k > 0$, and $-K_X$ thus admits a nonzero morphism in $kL = (kL + K_X) - K_X$, and so by composition with $\lambda^{\otimes k}$, $-K_X$ maps nontrivially into $\otimes^{mk} \Omega_X^1$. Equivalently, \mathcal{O}_X injects into $\otimes^{mk} \Omega_X^1 \otimes K_X$. Theorem 7.2 implies that K_X is pseudo-effective. \square

Remark 7.4. Let $X := \mathbb{P}^k \times Z$, where $k > 0$, and Z is of general type and dimension $(n - k) \geq 0$. Then K_X is not pseudo-effective, although $\Omega_X^{n-k} \subset \Omega_X^{1 \otimes (n-k)}$ contains $L := p^*(K_Z)$, a line bundle of Kodaira dimension $0 \leq (n - k) \leq (n - 1)$. The assumption that L is big cannot be weakened to any smaller Kodaira dimension in order to imply the pseudoeffectivity of K_X .

The argument used in Theorem 7.1 however extends to arbitrary numerical dimensions when K_X is assumed to be pseudo-effective.

Definition 7.5. Let L, P, A be line bundles on X , A ample.

1. $\kappa(X, L) := \max\{k \in \mathbb{Z} \mid \overline{\lim}_{m>0} (\frac{h^0(X, m.L)}{m^k}) > 0\} \in \{-\infty, 0, \dots, n\}$
2. $\nu(X, L) := \max\{A, k \in \mathbb{Z} \mid \overline{\lim}_{m>0} (\frac{h^0(X, m.L+A)}{m^k}) > 0\} \in \{-\infty, 0, \dots, n\}$

The elementary properties of these two invariants are:

Proposition 7.6. 0. $\nu(X, L) \geq \kappa(X, L)$

1. L is pseudo-effective (resp. \mathbb{Q} -effective) iff $\nu(X, L) \geq 0$ (resp. $\kappa(X, L) \geq 0$).
2. L is big (i.e. $\kappa(X, L) = n$) iff $\nu(X, L) = n$.
3. If P is pseudo-effective, $\nu(X, L + P) \geq \nu(X, L)$.
4. If L is big and if P is pseudo-effective, $L + P$ is big.

Theorem 7.7. Assume that K_X is pseudo-effective. Let L be a line bundle on X , and $L \subset \otimes^m \Omega_X^1$ a nonzero sheaf morphism. Then $\nu(X, L) \leq \nu(X, K_X)$.

Proof. Let $Q := (\otimes^m \Omega_X^1)/L^{sat}$. Since $\det(Q)$ is pseudo-effective by Theorem 7.1, $N.K_X = \det(L^{sat}) + \det(Q)$ is the sum of L^{sat} and $P := \det(Q)$ which is pseudo-effective.

Thus $\nu(X, K_X) = \nu(X, N.K_X) = \nu(X, L^{sat} + P) \geq \nu(X, L)$. \square

Remark 7.8. 1. The analogue of Proposition 7.6.(3) is false in general for the Kodaira dimension. The proof of Theorem 7.7 thus does not apply to prove $\kappa(X, L) \leq \kappa(X, K_X)$ in the same situation. However:

2. A central conjecture of birational geometry claims that $\nu(X, K_X) = \kappa(X, K_X)$ for any X . It thus implies, together with Theorem 7.7, that $\kappa(X, K_X) \geq \nu(X, L)$ for any $m > 0, L \subset \otimes^m \Omega_X^1$ if $\nu(X, K_X) \geq 0$.

3. The Remark 7.4 shows that the assumption that K_X is pseudo-effective cannot be removed or weakened.

When K_X is not pseudo-effective (i.e. when X is uniruled), we have the general version:

Corollary 7.9. *Let $f : X \rightarrow Z$ be the MRC of X . Let $L \subset \otimes^m \Omega_X^1$ be a subsheaf of rank 1. Then $\nu(X, L) \leq \nu(Z, K_Z)$.*

Proof. We can and shall assume that L is saturated in $\otimes^m \Omega_X^1 := E$. Let $X_z := f^{-1}(z)$, $z \in Z$ is a generic smooth fibre of f , and let A be an ample line bundle on X , with $L_{X_z} := L|_{X_z}$.

Lemma 7.10. $\nu(X, L) = -\infty$, unless $L_{X_z} \subset f^*(\otimes^m \Omega_{Z,z}^1)$ over X_z .

Proof. The exact sequence $0 \rightarrow f^*(\Omega_{Z,z}^1) \rightarrow \Omega_{X|X_z}^1 \rightarrow \Omega_{X_z}^1 \rightarrow 0$ induces on $\otimes^m \Omega_{X|X_z}^1$ a filtration with graded pieces isomorphic, for $0 \leq j \leq m$, to $G_j := \otimes^{m-j} f^*(\Omega_{Z,z}^1) \otimes (\otimes^j \Omega_{X_z}^1)$. Since X_z is rationally connected, we may choose $\alpha \in \text{Mov}(X)$ such that $f_*(\alpha) = 0$, and α restricts to X_z such that $\mu_{\alpha, \max}(\otimes^j \Omega_{X_z}^1) = j \cdot \mu_{\alpha, \max}(\Omega_{X_z}^1) < 0$, for any $j > 0$. If L is not contained in $\otimes^m (f^*(\Omega_{Z,z}^1))$, there exists $j > 0$ and an injective sheaf map $L \rightarrow G_j$. We thus get: $\mu_\alpha(L) = L \cdot \alpha \leq \mu_{\alpha, \max}(G_j) = j \cdot \mu_{\alpha, \max}(\Omega_{X_z}^1) < 0$, since $\mu_{\alpha, \max}(\otimes^{m-j} (f^*(\Omega_{Z,z}^1))) = 0$, the bundle $f^*(\Omega_{Z,z}^1)$ being trivial. Thus $L \cdot \alpha < 0$, and $\nu(X, L) = -\infty$. \square

From lemma 7.10, we deduce that $L \subset f^*(\otimes^m \Omega_Z^1)^{\text{sat}} \subset \otimes^m \Omega_X^1$. We can now conclude from the following Lemma 7.11, analogous to Lemma 7.1, using the very same arguments used to deduce Theorem 7.7 from Lemma 7.1.

Lemma 7.11. *Let $f : X \rightarrow Z$ be the MRC fibration of X , and let $\mathcal{G} \subsetneq f^*(\otimes^m \Omega_Z^1)^{\text{sat}}$ be a saturated subsheaf. Then $\det(\mathcal{Q})$ is pseudo-effective on X , where \mathcal{Q} is the quotient of $(f^*(\otimes^m \Omega_Z^1))^{\text{sat}}$ by \mathcal{G} .*

Proof. We may and shall assume Z to be smooth. By contradiction, if the conclusion does not hold, $\mu_{\alpha, \min}(f^*(\otimes^m \Omega_Z^1)^{\text{sat}}) < 0$ for some $\alpha \in \text{Mov}(X)$, and $-\mu_{\alpha, \max}((f^*(T_Z))^{\text{sat}}) = \mu_{\alpha, \min}((f^*(\Omega_Z^1)^{\text{sat}})) < 0$. There is then $\mathcal{F} \subset (f^*(T_Z))^{\text{sat}}$, α -semi-stable with $\mu_{\alpha, \min}(\mathcal{F}) > 0$ which is an algebraic foliation on X with closures of leaves rationally connected, and mapped non-trivially on Z by f , contradicting the non-uniruledness of Z . \square

\square

8. SHAFAREVICH-VIEHWEG CONJECTURE.

Let $f : Y \rightarrow B$ be a projective holomorphic submersion with connected fibres between connected complex quasi-projective manifolds Y, B .

We assume that all fibres of f have an ample canonical bundle, and that the ‘variation’ of f , defined by: $Var(f) = rk(ks : TB \rightarrow R^1 f_*(T_{Y/B}))$ is maximal, equal to $dim(B)$.

The situation considered by I. Shafarevich in 1962, was when f was a non-isotrivial family of curves of genus at least 2 on a curve B . His conjecture (proved by Parshin and Arakelov) was (formulated differently) that $K_{\bar{B}} + D$ was big on $\bar{B} = B \cup D$, a compactification of B .

The conjecture of Shafarevich was extended by Viehweg in the following form: in the above situation, $K_{\bar{B}} + D$ is big if \bar{B} is a smooth projective compactification of B obtained by adding to B a divisor D of simple normal crossings.

In this situation, Viehweg-Zuo ([37]) proved the existence of a big line bundle (the ‘Viehweg-Zuo sheaf’) $L \subset \otimes^m \Omega_{\bar{B}}^1(LogD)$ which, combined with the following Theorem, implies Viehweg’s conjecture (proved first in [32]):

Theorem 8.1. *Let X be a connected complex projective and D a divisor of simple normal crossings on X . Assume that a big line bundle L on X has an injective sheaf morphism $L \subset \otimes^m \Omega_X^1(LogD)$. Then $K_X + D$ is big.*

This is the logarithmic analogue of Theorem 7.3. It is proved by extending to the logarithmic setting the theorems 1.6 and 7.3, which is done very straightforwardly, by replacing in the proofs the (co)tangent bundles by their *Log*-analogs.

The Viehweg’s conjecture has been extended, and proofs given, in various directions (fibres of general type, or with semi-ample canonical bundle, or B ‘special’, and then f is isotrivial). In another direction, inspired by Lang’s conjectures, the Brody, Kobayashi, or Picard hyperbolicity of the corresponding moduli spaces have been proved in various situations. See [14] for references on this topic.

9. NUMERICALLY TRIVIAL FOLIATIONS.

Let $\mathcal{F} \subset TX$ be a numerically trivial and pseudo-effective foliation, that is: such that $det(\mathcal{F}).A^{n-1} = 0$, for some ample line bundle A . Since $c_1(\mathcal{F}) = 0$, the pseudo-effectivity of \mathcal{F} and \mathcal{F}^* are equivalent.

When X is an abelian variety and \mathcal{F} is linear (that is: $\mathcal{F} = Ker(w)$, $w \in H^0(X, \Omega^{n-r})$), \mathcal{F} is numerically flat and pseudo-effective, but the leaves of \mathcal{F} are in general not algebraic. In such cases however, \mathcal{F} is flat (i.e: given by a linear representation of the fundamental group), and so semi-stable but not stable.

The following question/conjecture was raised by Pereira-Touzot (who gave several affirmative answers, mostly in the corank-one case): if \mathcal{F} is A -stable and numerically trivial, is it algebraic?

Theorem 1.6, reduces this question to the non-pseudoeffectivity of \mathcal{F}^* when $c_1(\mathcal{F}) = 0$, and \mathcal{F} is A -stable.

A partial solution is given in the following:

Theorem 9.1. ([23],[12]) *Let E be a vector bundle on X , assume that $c_1(E) = 0 \neq c_2(E)$, and that $\text{Sym}^m(E)$ is A -stable for some ample A and some $m > r = \text{rk}(E)$. Then E is not pseudo-effective³.*

The proof of [12] relies on analytic and metric methods. The previous proof given in [23] by Höring-Peternell is mostly algebro-geometric. It runs along the following lines, starting with E pseudo-effective, with $c_1(E) = 0$, and all $\text{Sym}^m(E)$ A -stable, in order to show that $c_2(E) = 0$. The first step shows, by induction on the codimension of $W \subset P := \mathbb{P}(E)$, a component of the restricted base locus $B_-(L)$, $L := \mathcal{O}_P(1)$, that $p(W)$ has codimension at least 2 in X , $p : P \rightarrow X$ being the natural projection. The key ingredient in this step is following lemma, due to Mumford when $r = \text{rk}(E) = 2$:

Lemma 9.2. ([23]) *Let E be a vector bundle on a smooth projective curve C . Let $L := \mathcal{O}_{\mathbb{P}(E)}(1)$. Assume that $c_1(E) = 0$, and $\text{Sym}^m(E)$ is stable for all $m > 0$. Then $L^d \cdot Z > 0$ for all $d > 0$, and any irreducible $Z \subset \mathbb{P}(E)$ of codimension d .*

The second step then deduces from the first one that the restriction E_S of E to a general complete intersection surface $S \subset X$ of large degree, is nef. So that E and $\det(E^*) = \det(E) = 0$ are nef, and E^* is thus nef too. Hence E is numerically flat, so in particular all Chern classes of E vanish.

From Theorem 9.1 and Theorem 1.6, we get:

Corollary 9.3. *Let $\mathcal{F} \subset TX$ be a foliation with $c_1(\mathcal{F}) = 0 \neq c_2(\mathcal{F})$. If $\text{Sym}^{[m]}(\mathcal{F})$ is A -stable for some $m > \text{rk}(\mathcal{F})$, \mathcal{F} is algebraic.*

The Corollary 9.3 (or the slightly weaker version of [23]) now permits to give an alternative proof of the Beauville-Bogomolov-Yau⁴ (BBY) decomposition theorem in the projective and klt singular case.

10. THE BEAUVILLE-BOGOMOLOV-YAU DECOMPOSITION

Let X be an n -dimensional compact Kähler manifold with $c_1(X) = 0$. By Yau's theorem, any such X carries a Ricci-flat Kähler metric. There are 3 basic examples of such manifolds, each class characterised by the restricted holonomy group Hol^0 of any of their Kähler, Ricci-flat, metrics: Hol^0 is trivial for 1, is $Sp(4r)$ for 2, and is $SU(n)$ for 3.

1. The compact complex tori are the quotients \mathbb{C}^n/Γ , where Γ is a cocompact lattice of \mathbb{C}^n .

³The proof works for X compact Kähler, and E a reflexive sheaf on X .

⁴Usually called the Bogomolov-Beauville decomposition. Since Yau's Ricci-flat metric is essential here, BBY seems justified.

2. The irreducible hyperkähler (HK) manifolds are defined as the ones which are simply-connected, even-dimensional: $n = 2r$, and admit a holomorphic symplectic 2-form σ generating $H^0(X, \Omega_X^2)$, and such that $s^{\wedge r}$ is a nowhere vanishing section of K_X . There are few known examples: 2 deformation classes in each even dimension due to Beauville (and Fujiki in dimension 4), and one additional in dimensions 6 and 10, found by O’Grady.

3. The Calabi-Yau (CY) manifolds are those which are simply connected, with $h^{p,0} = 0, \forall p = 1, \dots, (n - 1)$, and with K_X trivial. Many deformation families known in each dimension. Among these complete intersections of suitable multidegrees in the projective spaces.

The BBY decomposition is the following:

Theorem 10.1. ([2]) *Let X be a connected compact Kähler manifold with $c_1(X) = 0$. A suitable finite étale cover of X is a product of a complex compact torus T , and of simply-connected CY or HK manifolds.*

Let us sketch the classical proof given in [2]: We equip X with some Ricci-flat Kähler metric ([38]). Let Hol^0 be its restricted holonomy representation, and $T_X = F \oplus (\oplus_i T_i)$ be the splitting of the tangent bundle of X into factors which are irreducible for the action of Hol^0 . This splitting is well-defined locally on X , and globally only up to possible reordering. These local factors correspond also to a local splitting of X into a direct product of Kähler submanifolds. By Berger general classification of the holonomy representations, only a flat factor F appears, together with other factors T_i having restricted holonomy either Sp or SU (the ones susceptible to preserve a section of K_X by parallel transport).

We lift this decomposition to the universal cover \tilde{X} of X on which the metric is complete. By De Rham decomposition theorem, \tilde{X} splits as a Kähler product, corresponding to these Hol^0 -factors ([3, Theorem 10.43]). The Cheeger-Gromoll theorem then says that $\tilde{X} = \mathbb{C}^k \times P$, where \mathbb{C}^k corresponds the flat factor, the other factor P being compact, and corresponding to the product of the factors with Hol^0 either Sp or SU ([3, Theorem 6.65]). Bieberbach’s theorem ([4]) then concludes the proof.

Notice that this shows in particular that the (everywhere regular) foliations defined on X by the factors T_i have compact leaves (with finite fundamental groups), and are thus ‘algebraic’ (i.e: with equal Zariski and topological closures of their leaves).

We shall now give, but only in the projective case, an alternative proof which does not require the consideration of the universal cover. This is done by proving directly, using Theorem 9.1, that the foliations defined by the T_i ’s are all algebraic. This follows from the following:

Lemma 10.2. *For each i , and each $m > 0$, $c_1(T_i) = 0$, and $Sym^m(T_i)$ is A -stable for any ample line bundle A on X .*

Proof. Since $c_1(TX) = c_1(F) = 0$, the first claim follows from the second one. The second claim is a consequence of representation theory of the groups SU and Sp ([17]). \square

If we take thus the (regular and submersive) fibration $f_i : X \rightarrow B$ with fibres the (projective) leaves of T_i , it is isotrivial of fibre F_i (by the existence of the transversal foliation defined by $F \oplus_{j \neq i} T_j$), and F_i has a vanishing irregularity and a discrete automorphism group (because the restricted holonomy is either SU or Sp), which implies that after a finite étale base change $B' \rightarrow B$, X splits as a product $F_i \times B'$.

Induction on $\dim(X)$ now reduces the proof to the case where $TX = F$, the flat factor. Bieberbach theorem ([4]) then permits to conclude. For details, see [9].

This BBY decomposition is still valid for projective varieties with *klt* singularities, but the proof, obtained first in [23], and using a combination of the works [16], [22], [20], [15], is much more involved, both technically and conceptually.

A simplified proof, which closely follows the above alternative proof in the smooth case, and avoids the delicate arguments in positive characteristic used in [15], can be found in [9]. The result has then been extended to the Kähler case in [1], by deformation to the projective case.

11. SOME QUESTIONS

We denote by X a smooth and connected complex projective manifold⁵, of dimension n . Let \mathcal{E} be a torsionfree coherent sheaf on X .

11.1. Pseudoeffectivity of determinants.

Definition 11.1. *We say that \mathcal{E} is pseudo-effective if so is $\det(Q)$, for each nonzero quotient of \mathcal{E}^{**} . Equivalently, $\mu_{\alpha, \min}(\mathcal{E}^{**}) \geq 0, \forall \alpha \in \text{Mov}(X)$.*

Example 11.2. *1. If \mathcal{E} is pseudo-effective, so is any of its quotients, as well as $\otimes^m \mathcal{E}, Sym^m \mathcal{E}, \wedge^m \mathcal{E}, \forall m > 0$, and more generally the tensors deduced from \mathcal{E} by linear representations, by Theorem 4.6.*

2. If $K_X = \det(\Omega_X^1)$ is pseudo-effective, so is Ω_X^1 , as seen in Theorem 7.1. In [10], the same property is shown, more generally, for any quotient of $\otimes^m(\Omega_X^1)$ with pseudo-effective determinant.

We ask the same question also for subsheaves of $\otimes^m(\Omega_X^1)$:

⁵The questions raised below could be extended to the compact Kähler case, and to smooth or klt orbifold pairs.

Question 11.3. *Let $\mathcal{E} \subset \otimes^m \Omega_X^1$ be such that $\det(\mathcal{E})$ is pseudo-effective. Is \mathcal{E} pseudo-effective? This is a birational version of stability.*

Notice that if $\mathcal{E} \subset \Omega_X^p$ is saturated and defines an *algebraic* foliation of rank $r = (n - p)$, the answer is yes by [10], since \mathcal{E} is then the cotangent sheaf of the space of leaves.

This question might be answered by showing the relative algebraicity of two transcendental foliations deduced from suitable quotients of Ω_X^1 , using a suitable extension of the next algebraicity criterion in [10].

Recall this algebraicity criterion (theorem 1.7): if $\mathcal{F} \subset TX$ is a foliation such that \mathcal{F}^* (which is a quotient of Ω_X^1) is not pseudo-effective, the leaves of \mathcal{F} are algebraic. Moreover, if $\mu_{\alpha, \max}(\mathcal{F}^*) < 0$ for some $\alpha \in \text{Mov}(X)$, the leaves of \mathcal{F} are rationally connected.

Question 11.4. *If \mathcal{F}^* is not pseudo-effective, are there restrictions on the leaves of \mathcal{F} , and which ones?*

Already when $\mathcal{F} = TX$, the answer is nontrivial, and consists in the classification of manifolds X with Ω_X^1 not pseudo-effective, or equivalently with $\nu_1(X) = -\infty$, using the notations of the next subsection. Although such manifolds may have an ample, or a trivial canonical bundle, as shown by hypersurfaces in \mathbb{P}^{n+1} , they share some common properties with the rationally connected manifolds. For example, rationally connected manifolds are simply-connected, and it is shown in [8] that the linear representations of X have a finite image if Ω_X^1 is not pseudo-effective. For manifolds of general type, even for surfaces, the invariant κ_1 (see below), does not seem to be described by, or even related to, other algebro-geometric invariants. For special manifolds, in the sense of definition 11.11 below, the situation might be much simpler (see conjecture 11.13).

An additional problem is that if the foliation \mathcal{F} has \mathcal{F}^* not pseudo-effective, and so if \mathcal{F} is algebraic, it is not clear whether or not the general fibres of an associated fibration have Ω^1 not pseudo-effective. The proof when $\mu_{\alpha, \min}(\mathcal{F}) > 0$ relies on the MRC, and it is unknown whether a similar fibration may be expected to exist on any X , with fibres having Ω^1 not pseudo-effective, and base with Ω^1 pseudo-effective.

11.2. Variations on Abundance. Let \mathcal{E} be torsionfree coherent on X , of rank r , and let $\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathcal{E}}(1)$ be the corresponding (Grothendieck) projectivisation, together with its tautological line bundle.

Definition 11.5. *We write $\kappa(\mathcal{E}) := \kappa(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathcal{E}}(1)) - (r - 1)$, and $\nu(\mathcal{E}) := \nu(\mathbb{P}(\mathcal{E}), \mathcal{O}_{\mathcal{E}}(1)) - (r - 1)$. We also write $\nu_p(X) := \nu(\Omega_X^p)$, and $\kappa_p(X) := \kappa(\Omega_X^p)$, so that: $\nu(X) = \nu_n(X)$, and $\kappa(X) := \kappa_n(X)$.*

The definition fits with the original one for line bundles. Moreover, the invariants κ_p and ν_p are preserved under finite étale covers. The invariant κ_1 has been introduced and studied by F. Sakai in [36]⁶.

The classical ‘Abundance conjecture’ $\nu_n(X) = \kappa_n(X)$ extends as:

Conjecture 11.6. *For any X and $p > 0$, one has: $\nu_p(X) = \kappa_p(X)$.*

Remark 11.7. *1. In [24], this is stated when $\kappa_p(X) = -\infty$, and proved for elliptic surfaces (except for some isotrivial ones), and some other additional cases.*

2. If $\kappa_p(X) = -\infty, \forall p > 0$, and if we assume the classical ‘Abundance conjecture’, we get by the MRC fibration that X is rationally connected, and so that $-\infty = \kappa_p(X) = \nu_p(X), \forall p > 0$. Conjecture 11.6 is thus a consequence of its ‘classical’ version in this situation.

Example 11.8. *1. If $c_1(X) = 0$, from the Beauville-Bogomolov-Yau decomposition theorem, and [23], one can easily see⁷ that $\nu_1(X) = \kappa_1(X) = \tilde{q}(X) - n$, where $\tilde{q}(X)$ is the maximum of $q(X')$ when X' runs through the finite étale covers of X . The same properties can be still shown with more work⁸ when X is a smooth model of a klt variety with $c_1 = 0$. Thus $\kappa_1(X) = \nu_1(X) = \tilde{q}(X) - n$ whenever $\kappa(X) = 0$, if one assumes the existence of good minimal models.*

Remark 11.9. *The Abundance conjecture $\nu = \kappa$ fails in general for rank-one subsheaves L of Ω_X^p , by the examples of M. Brunella and M. Mc Quillan of the (duals of the) tautological foliations on irreducible free quotients of the bidisc, which have $\kappa = -\infty$ and $\nu = 1$.*

Question 11.10. *Are there rank-one subsheaves L of Ω_X^p for suitable $X, p > 0$, with $p = \nu(X, L) > \kappa(X, L) \geq 0$? As seen just above, there are examples with $p = \nu(X, L) > \kappa(X, L) = -\infty$.*

11.3. Special manifolds. Recall that from Bogomolov’s theorem that $\kappa(X, L) \leq p, \forall X, \forall p > 0$, if L is any rank-one subsheaf of Ω_X^p .

Definition 11.11. *We say that X is special⁹ if $\kappa(X, L) < p$, for any such L , and any $p > 0$;*

Bogomolov’s theorem has been strengthened by C. Mourougane and S. Boucksom to $\nu(X, L) \leq p, \forall X, \forall p > 0$.

Conjecture 11.12. *If X is special, $\nu(X, L) < p, \forall L \subset \Omega_X^p, \forall p > 0$, L of rank one.*

⁶F. Sakai uses the ‘invariant λ ’, which takes the value $-n$ when $\kappa_1 = -\infty$, but otherwise coincides with κ_1 . His convention simplifies the formulations in some situations, such as products.

⁷In the next formula, if $\tilde{q}(X) = 0$, replace Sakai’s $-n$ by $-\infty$.

⁸Because one needs to carefully distinguish étale from quasi-étale covers, as shown by the Kummer $K3$ ’s.

⁹We may assume X to be compact Kähler, here.

This has been proved in [34] for $p = 1$. Notice that if $L \subset \Omega_X^1$ is of rank one with $\nu(X, L) = 1$, there may be no $L' \subset \Omega_X^1$ with $\kappa(X, L') = 1$, as shown by the example of Brunella and Mc Quillan in remark 11.9, since no finite étale cover of the ambient surface maps onto a curve of positive genus.

Conjecture 11.13. *Assume X (smooth compact Kähler) is special. Do we have¹⁰ : $\nu_1(X) = \kappa_1(X) = \tilde{q}(X) - n$?*

If yes, then $\nu_1(X) = \kappa_1(X) = -\infty$, that is Ω_X^1 not pseudo-effective, if and only if $\tilde{q}(X) = 0$, if and only if $\pi_1(X)$ is finite.

Recall that the Abelianity conjecture claims that $\pi_1(X)$ is virtually abelian if X is special, so that $\tilde{q}(X) = 0$ should then mean that $\pi_1(X)$ is finite.

Notice that if X is a compact Kähler complex manifold with all of its finite étale covers having a surjective Albanese map (as is the case when X is special), then $\kappa_1(X) \geq \tilde{q}(X) - n$.

Acknowledgement: We thank the reviewers for their careful readings, corrections and suggestions, which lead to improvements of the text.

REFERENCES

- [1] Bakker-H. Guenancia-C. Lehn. Algebraic approximation and the decomposition theorem for Kähler Calabi-Yau varieties. arXiv. 2012.00441
- [2] A. Beauville. Variétés Kählériennes dont la première classe de Chern est nulle. J. Diff. Geom. 18, 755-782 (1983)
- [3] A.L. Besse. Einstein manifolds. Ergeb.Math.Grenz.Geb.10. Springer Verlag (1987)
- [4] L. Bieberbach. Über die Bewegungsgruppen des Euklidischen Raumes. I,II. Math. Ann. 70, 297-336 (1911), 72, 400-412 (1912)
- [5] F. Bogomolov. Hamiltonian Kähler manifolds. Sov. Math. Dokl. 19, 1462-1465 (1978).
- [6] F. Bogomolov-M. McQuillan. Rational curves on foliated varieties. IHES/M/01/07 (2001).
- [7] S. Boucksom-JP. Demailly-M. Păun-T. Peternell. The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension. J. Alg. Geom. 22 (2013), 201-248.
- [8] Y. Brunella-B. Klingler-B. Totaro. Symmetric differentials and the fundamental group. Duke Math. J. 162 (3013), 2797-2813.
- [9] F. Campana. The Bogomolov-Beauville-Yau decomposition for KLT Projective Varieties with trivial first Chern class-without tears. Bull. SMF. 149 (2021), 1-13.
- [10] F. Campana-M. Păun. Foliations with positive movable slope. Publ. Math. IHÉS vol. 219, p. 1-49 (2019)
- [11] F. Campana-T. Peternell. Geometric stability of the cotangent bundle and the universal cover of a projective manifold. Bull. SMF (2011)
- [12] F. Campana-J. Cao-M. Păun. Subharmonicity of direct images and applications. ArXiv 2019.

¹⁰If $\tilde{q}(X) = 0$, replace the RHS $-n$ by $-\infty$.

- [13] B. Claudon. Positivité du cotangent logarithmique et conjecture de Shafarevich-Viehweg. Séminaire Bourbaki. Exposé n^0 1105. Novembre 2015.
- [14] Y. Deng. Big Picard theorem and algebraic hyperbolicity for varieties admitting a variation of Hodge structures. arXiv:2001.04426
- [15] S. Druel. A decomposition theorem for singular spaces with trivial canonical class of dimension at most five. *Inv. math.* 211, 245-296 (2018)
- [16] P. Eyssidieux-V. Guej-A. Zeriahi. Singular Kähler-Einstein metrics. *J. AMS.* 22 (2009), 607-639.
- [17] W. Fulton-J. Harris. Representation theory. A first course. Springer-Verlag (1991).
- [18] F. Gounelas. Free curves on varieties. *Doc. Math.* 21 (2016), 287-308.
- [19] T. Graber-J. Harris-J. Starr. Families of rationally connected varieties. *J. Amer. Math. Soc.* 16 (2003), p. 57-67.
- [20] D. Greb-H. Guenancia-S. Kebekus. Klt varieties with trivial canonical class, holonomy, differential forms, and fundamental groups. *Geometry and Topology* 23-4 (2019), 2051-2124.
- [21] D. Greb-S. Kebekus-T. Peternell. Movable curves and semistable sheaves. *IMRN* (2016), 536-570.
- [22] D. Greb-S. Kebekus-T. Peternell. Etale fundamental group of Kawamata log-terminal spaces, flat sheaves and quotients of abelian varieties. *Duke Math. J.* Volume 165, Number 10 (2016), 1965-2004.
- [23] A. Höring-T. Peternell. Algebraic integrability of foliations with numerically trivial canonical bundle. *Inv. Math.* 216, 395-419 (2019)
- [24] A. Höring-T. Peternell. A nonvanishing conjecture for cotangent bundles. arXiv 2006. 05225.
- [25] S. Kobayashi. Differential geometry of vector bundles. *Publ. Math. Soc. Japan.* Vol. 15 (1987)
- [26] J.Li-S.T.Yau. Hermitian Yang-Mills connections on non-Kähler manifolds, 560-573, in *Mathematical aspects of String theory.* S.T. Yau Ed; World Scientific Publish. London (1987).
- [27] M. Lübke-A. Teleman. The Kobayashi-Hitchin correspondance. World Sc. Publish. (1995).
- [28] Y. Miyaoka. Deformations of a morphism along a foliation and applications, in *Algebraic geometry Bowdoin* (1985), *Proc. Symp. Pure Math.* Vol. 46 (1987), p. 245-268.
- [29] Y. Miyaoka- S. Mori. A numerical criterion for uniruledness. *Ann. Math.* 124 (1986), 65-69.
- [30] D. Mumford. Hilbert fourteenth problem-the infinite generation of subrings such as rings of invariants. *Proceedings of Symposia in pure mathematics.* Vol 28 (1976), 431-443.
- [31] N. Nakayama. Zariski decomposition and Abundance. *MSJ Memoirs*, Volume 14 Tokyo, The Mathematical Society of Japan, 2004
- [32] B. Taji. The isotriviality of smooth families of canonically polarized manifolds over a special quasi-projective base. *Comp. Math.* Vol. 152 (2016), 1121-1134.
- [33] J. Ottem. On subvarieties with ample normal bundle. *J. Eur. Math. Soc.* vol.18 (2016), 2459-2468.
- [34] J. Pereira-E.Rousseau-F.Touzet. Numerically nonspecial varieties. arXiv. 2106.12275.
- [35] T. Peternell. Compact subvarieties with ample normal bundles, algebraicity, and cones of cycles. *Mich. Math. J.* Vol. 61 (2012), 875-889.
- [36] F. Sakai. Symmetric powers of the cotangent bundle and classification of algebraic varieties. *LNM* 732, 545-563.

- [37] E. Viehweg-K. Zuo. Base spaces of non-isotrivial families of smooth minimal models, in Complex geometry (Göttingen 2000). Springer-Verlag (2002), p. 279-328.
- [38] S.T. Yau. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation I, Comm. Pure and Appl. Math. 31, 339-411 (1978)

UNIVERSITÉ LORRAINE, NANCY,

E-mail address: `frederic.campana@univ-lorraine.fr`